

## Assay system to screen for compounds inducing PR-gene expression in grape vine (*Vitis spec.*)

Tobias Seibicke<sup>1</sup>, Alexander Rügner, Gunther Neuhaus<sup>2</sup>,  
Hanns-Heinz Kassemeyer, Günther Buchholz

### Abstract

Plants respond to pathogen attack by activation of an array of inducible defense responses e.g. PR-proteins (pathogenesis related). Transcription of a  $\beta$ 1,3 glucanase gene was found to be induced in grape leaves upon infection with *Plasmopara viticola*. Promoter- and coding sequences of the gene were identified. In transient transformation assays the glucanase::GFP-fusion product was found to be localized exclusively in the cytoplasm of protoplasts derived from a *Vitis* cell culture. A fusion of the glucanase-promoter and the firefly luciferase-coding region was found to be induced by addition of salicylic acid (SA) in transiently transformed protoplasts. The same construct was used to create a stably transformed *Vitis* suspension cell culture for screening and detecting compounds acting as elicitors effectively inducing this PR promoter.

### Keywords

PR-proteins, glucanase promoter, luciferase, *Vitis*, plant cell culture, elicitor

---

<sup>1</sup> Staatliches Weinbauinstitut Freiburg, Merzhauser-Str. 119 D-79100 Freiburg (Germany)

<sup>2</sup> Institut für Biologie II (Zellbiologie), Schänzlestr. 1, D-79104 Freiburg (Germany)